

## Predator Diversity of Mayflies from Kolhapur District, India

Rohini Kamble<sup>1</sup> and T.V.Sathe<sup>2\*</sup>

<sup>1-3</sup> Dept. of Zoology, Shivaji University, Kolhapur– 416004, India

Email: : [profdrtvsathe@rediffmail.com](mailto:profdrtvsathe@rediffmail.com); [rohinistr36@gmail.com](mailto:rohinistr36@gmail.com)

### ABSTRACT

Mayflies (Ephemeroptera) are the most primitive and ancient insect group globally; there are about 3000 species of mayflies belonging to 400 genera and 42 families. Out of which 390 species belonging to 84 genera and 20 families occur in the oriental region. From India, 124 species of mayflies have been recorded under 46 genera and 12 families. Adult mayflies are terrestrial and immature forms are aquatic and called naids (nymphs). Mayflies are good source for food for various animal groups such as Insects, Platyhelminthes, Spiders, Pisces, Amphibians and Aves. Therefore, biodiversity of predatory species of mayflies have been studied as nutrition affect the life cycle, survival, fecundity, sex ratio, etc. A list of 40 predatory species of above said groups has been reported from rivers Panchganga, Krishna, Warna, Bhogavati, Kumbhi, Kasari, and Saraswati of Kolhapur district.

**Key words** – Predator diversity, Mayflies, Rivers, Kolhapur district

### INTRODUCTION

The nutritive value of food significantly affects the life history characteristics of a variety of aquatic insect species (Sweeney, 1984). Several studies have focused on food as an experimental variable and the interaction between temperature and food in aquatic ecosystem (Anderson & Cummins, 1979). For aquatic detritivores, studies of food quality have largely involved rearing larvae for various length of time on different species of deciduous leaves (Wallace *et al.*, 1970). Leaf species often differ from one another in terms of biochemical composition and associated microflora. Larval feeding and growth rates are vary significantly with the species of leaf provided as food (Cummins *et al.*, 1973).

Herbivory is an important factor regulating the biomass and community structure in streams (Feminella and Howkins, 1995). Stream food webs rely primarily on two food sources: autochthonous primary production within the stream and autochthonous organic matter

transferred to the stream as leaf litter, woody debris and dissolved organic carbon. Autochthonous algae are more important than autochthonous organic matters to stream consumers (March and Pringle 2003) since they are the main energy source in some headwater streams (Mantel *et al.*, 2004, Lau *et al.*, 2008, 2009, Li and Dudgeon 2008). Considering the fact that Kolhapur district abounds in tanks, ponds, small lake reservoirs and wetlands of various sizes, the diversity of mayflies and their predators will add great relevance in sustainable development of this region. Mayflies (Insecta: Ephemeroptera) are pollution indicators. Therefore, it is necessary to investigate their mortality factors in terms of their predation.

Review of literature indicates that mayflies have been studied by several workers (Fladung, 1924; Bodenheimer, 1951; Martof & Scott, 1957; Wallace *et al.*, 1970; Cummins *et al.*, 1973; Downes, 1978; Brittain, 1978; Anderson & Cummins, 1979; Reinecke *et al.*, 1980; Soluk & Clifford, 1984; Sweeney, 1984; Bart & Holzenthal, 1985; Nost, 1985; Gibbs & Mingo, 1986; Giller, 1986; Scrimgeour & Winterbourn, 1987; Bergeron *et al.*, 1988; Dudgeon, 1989; Harries, 1990; Ormerod & Tyler, 1991; Kjellberg, 1992; Rabeni, 1992; Gupta *et al.*, 1993; Caira *et al.*, 1995; Castien & Gosalbez, 1995; March & Pringle, 2003; Mantel *et al.*, 2004; Lau *et al.*, 2008, 2009; Li & Dudgeon, 2008; Sathe, 2014, 2015), Kamble & Sathe, 2015, 2016; Sathe & Patil, 2016) from the review of their predation and seasonal abundance from different part of the world.

#### How to Site This Article:

Rohini Kamble and T.V.Sathe (2017). Predator Diversity of Mayflies from Kolhapur District, India. *Biolife*. 5(1), pp 85-91

DOI: 10.5281/zenodo.7360284

Received: 6 January 2017;

Accepted: 27 February 2017;

Available online : 5 March 2017

## MATERIALS AND METHODS

### Study site:

Diversity of mayflies specially their nymph and adults and their predators both from vertebrates and invertebrates have been studied from rivers in Kolhapur district. (Lat.  $16^{\circ} 41' 44''$  N, Long  $74^{\circ} 13' 54''$  E Alt. 184 F, 543msl). Kolhapur district (Fig.1) receives an annual rainfall of about 1138.5 mm, most of which occurs during June to October.

Figure-1. Rainfall of Kolhapur district



Collections were made from marginal weed beds of rivers beside, some predators have been dissected which were collected from various rivers viz,

- I. Krishna - Shirol (Alt-1732 F, 527msl, Lat-  $16^{\circ} 61' N$ , Long-  $74^{\circ} 36' E$ ),
- II. Panchganga – Karveer (Alt-1867 F, 543msl, Lat-  $16^{\circ} 41' N$ , Long-  $74^{\circ} 13' E$ ),
- III. Warna - Hatkangale (Alt -1939 F, 591msl, Lat-  $16^{\circ} 61' N$ , Long-  $74^{\circ} 35' E$ ),
- IV. iv)Kasari - Panhala (Alt-2903 F, 845msl, Lat-  $16^{\circ} 82' N$ , Long-  $74^{\circ} 12' E$ ),
- V. Kasari - Shahuwadi (Alt-1903 F, 580msl, Lat-  $16^{\circ} 85' N$ , Long-  $73^{\circ} 90' E$ ),
- VI. vi)Bhogavati - Radhanagri (Alt -2034 F, 619msl, Lat-  $16^{\circ} 24.50' N$ , Long-  $74^{\circ} 59.52' E$ ),
- VII. vii) Kumbhi - Gaganbavda (3177 F, 900msl, Lat-  $16^{\circ} 69' N$ , Long-  $74^{\circ} 107' E$ ),
- VIII. viii) Sarasvati - Gaganbavda (3177 F, 900msl, Lat-  $16^{\circ} 69' N$ , Long-  $74^{\circ} 107' E$ ) and observations have been made on the availability of body parts of mayflies in the gut of predators. The selection of study spots was based on variations in rainfall and altitude.

### Experimental set up:

Air and water temperature readings taken with a mercury bulb thermometer at the time of collection of water and insect samples. Nymphs of Mayflies species and the other aquatic predatory insects collected by

dragging a 25 x 25 cm net of 200  $\mu$ m mesh through a 2m long section of the weed bed and density finally expressed as numbers per unit sampling effort. Three replicate samples collected at fortnightly interval. Nymphs preserved in the field, in 6% formalin to prevent possible digestion of food materials and sorted in the laboratory under a dissecting binocular microscope. Body length (excluding antenna and cerci) and head width measured with a calibrated ocular micrometer. Mayflies emergence data were obtained by collecting submagines from the underside of leaves of the lake side vegetation and by rearing nymphs in the laboratory. The foregut contents of predatory nymphs/ larvae/ adult pooled and mixed with water of known volume in a tube. This suspension was then vigorously agitated and 1 ml subsample transferred to a small counting cell fitted on a glass slide and examined under a microscope equipped with a squared ocular grid.

Five to ten such subsamples examined. The number of diatoms, desmids, flagellates, and cells of filamentous algae in each subsample as well as the number of squares in the grid occupied by detritus and mineral particles counted. (Shapas & Hisenhoff 976). The predators identified as feeding on mayflies by gut or fecal analysis or by direct observation of feeding.

## RESULTS

Diversity of Mayflies and their predators recorded in table-1 and figs- 1 to 11 indicated that a total no. of 40 species of predators from invertebrates and vertebrates were prevalent in Kolhapur region. While, the mayflies predated were from more than ten species (table-1). The most abundant of mayflies recorded were *Baetis* spp., *Leptophelebia* sp., *Heptanigids* and *Indialis* sp. Desmids and filamentous green algae were abundant during May to October, especially July to September while, detritus gained importance food during November to March. Flagellates were abundant during November to April. Diatoms were found in the gut almost throughout the year.

Population size of mayflies was relatively small during December to February 2015-2016. Thereafter, from October onward, density built up steadily to reach a peak in May to June after which it declined, although a sizeable population persisted in June. Reduced density in July to October was most likely to have been caused by low temperature rains and flooding in Kolhapur District.

## DISCUSSION

According to Gibbs and Mingo (1986), some mayflies are predators of other mayflies. *Siphonisca aerodromia* nymph begin their lives as small detritivores but these agile, rapid swimmers became more carnivorous as the age, feeding on the likes of *Siphonurus*, *Leptophelebia* and *Ephemerella*. The sand dwelling heptageniid, *Pseudiron centralis*, preferred chironomids as food, 5%

Table-1. Predators of Mayflies from Kolhapur District

| Sr. No                                  | Predators   | Mayflies   | Prey stage attacked   | Distribution  |
|---|---|--|---|---|
| 1.<br>2.                                | <b>Invertebrates :</b><br>Platyhelminthes :<br><i>Tricladida</i> sp.<br><i>Dugecia</i> sp.  | <i>Baetis</i> sp.<br><i>Baetis</i> sp.   | Nymph<br>Nymph  | AJR, RDR<br>SR, KR, HTK                                     |
| 3.<br>4                                 | Arthropoda :<br>Ephemeroptera :<br><i>Ephemerella</i> sp.<br><i>Baetis</i> sp.  | <i>Indialis badia</i><br><i>Indialis badia</i>   | Nymph<br>Nymph  | HTK, KR<br>HTK  |
| 5<br>6<br>7<br>8<br>9<br>10<br>11<br>12 | Odonata :<br><i>Libellula depressa</i><br><i>Anex</i> sp.<br><i>Pantala</i> sp.<br><i>Crocothemis survilia</i><br><i>Ischnura</i> sp.<br><i>Ceragrion</i> sp.<br><i>Lestes</i> sp.<br><i>Penueus indicus</i>                      | <i>Leptophelebia</i> sp.<br><i>Ephemerella</i> sp.<br><i>Indialis badia</i><br><i>Caenis</i><br><i>Leptophelebia</i> sp.<br><i>Baetis</i> sp.<br><i>Baetis</i> sp<br><i>Caenis</i> | Nymph, Adult<br>Nymph, Adult<br>Nymph, Adult<br>Nymph, Adult<br>Nymph, Adult<br>Nymph, Adult<br>Nymph, Adult<br>Nymph | KR, RDR<br>GGD<br>KR, HTK<br>PNH<br>PNH<br>SHW<br>HTK<br>KR |
| 13<br>14<br>15                          | Hemiptera :<br><i>Ranatra elongata</i><br>Scorpion bug<br><i>Notonecta</i> sp.  | <i>Indialis badia</i><br><i>Baetis</i> sp.<br><i>Leptophelebia</i> sp.   | Nymph<br>Nymph<br>Nymph   | KR, GGB, RDR<br>PNH   |
| 16<br>17<br>18                          | Acarina : Spiders<br><i>Lycosa</i> sp.<br><i>Hippasa</i> sp.<br><i>Pandosa</i> sp.  | Heptaginids<br>Heptaginids<br>Heptaginids  | Adult<br>Adult<br>Adult   | KR,GGB<br>RDR   |
| 19<br>20<br>21<br>22<br>23<br>24<br>25  | <b>Vertebrates : Pisces :</b><br><i>Poecilla reticulata</i><br><i>Cyprinus</i> sp.<br><i>Oreochromis</i><br><i>mossambica</i><br><i>Gambusia affinis</i><br><i>Clarias batrochus</i><br><i>Labeo rohita</i><br><i>Catla catla</i> | <i>Baetis</i> sp.<br><i>Caenis</i> sp.<br><i>Indialis badia</i><br><i>Leptophelebia</i> sp.<br><i>Baetis</i> sp.<br><i>Indialis badia</i><br><i>Indialis badia</i>                 | Nymph<br>Nymph<br>Nymph<br>Nymph<br>Nymph<br>Nymph, Adult<br>Nymph  | SR, AJR<br>GGD<br>SR<br>PNH<br>KR, HTK<br>SHW<br>PNH        |
| 26<br>27<br>28<br>29<br>30              | Amphibia :<br><i>Bufo</i> spp.<br><i>Rana</i> sp.<br><i>Duttaphrynus</i> sp.<br><i>Euphyetis</i> sp.<br><i>Polypedatus</i> sp.  | <i>Caenis</i> sp.<br><i>Baetis</i> sp.<br><i>Indialis</i> sp.<br><i>Indialis</i> sp.<br><i>Indialis</i> sp.  | Adult<br>Nymph, Adult<br>Adult<br>Nymph, Adult<br>Nymph, Adult  | KR<br>RDR<br>SR<br>GGD<br>SR                                |
| 31<br>32                                | Reptalia :<br><i>Salamandra</i> sp.<br><i>Euphaea decorata</i>  | <i>Leptophelebia</i> sp.<br><i>Ephemerella</i> sp.   | Adult<br>Adult  | GGB<br>SR, HTK  |
| 33<br>34<br>35<br>36<br>37              | Aves :<br><i>Merops orientalis</i><br><i>Amauornis phoenicurus</i><br><i>Ardeola groyii</i><br><i>Pscnonotus cafer</i><br>Wagtail   | <i>Baetis</i> sp.<br><i>Leptophelebia</i> sp.<br><i>Leptophelebia</i> sp.<br><i>Baetis</i> sp.<br><i>Indialis badia</i>  | Adult<br>Nymph, Adult<br>Nymph<br>Adult<br>Adult  | KR<br>CGD,KR<br>SR<br>RDR<br>PNH                            |
| 38<br>39<br>40                          | Mammalia : Chiroptera :<br><i>Pteropus giganteus</i><br><i>Hipposideros speoris</i><br>Primates :<br><i>Homo sapiens</i>  | Heptaginids<br>Heptaginids<br>Heptaginids  | Adult<br>Adult<br>Nymph, Adult  | SR, KR<br>PNH<br>RDR  |

Note – SR- Shirol, KR – Karveer, HTK – Hatkangale, PNH – Panhala, SHW – Sahuwadi, GGD – Gaganbavda, RDR – Radhanaqri.



Figure-2. *Indialis Badia* Adult



Figure-3. *I. badia* Nymph



Figure-4. *Ephemerella* Nymph



Figure-5. *Ischunura* sp. Adult



Figure-6. Damselfly Nymph



Figure-7 *Crocothemis* Nymph




Figure-8. *Ranatra* sp. AdultFigure-9. *Lycosa* sp.Figure-10. *P. indicus* sp

Figure-11. Frog tadpole



of its foregut contents was composed of early instars of *Baetis*, *Centroptilum* and *Ephron* (Soluk and Clifford, 1984). Agnew (1962), found remains of baetid nymphs in two of seven guts in *Centroptiloides bifasciata* and *Ephemerella* sp. In the nymphs of above two species 50%, gut was fitted with *Hetagenia* nymphs (Muttkowski and Smith, 1929).

Downes (1978) has observed six species of Ceratopogonids (Diptera) feeding on mayflies. The female Ceratopogonid typically entered in to the mayfly swarm and landed on the thorax for predation. Fifty species of Podagritis wasps (Hymenoptera: Sphecidae) were distributed throughout America, Australia and New Zealand as dipterivorous (Harris, 1990). Harries (1990) reported two species *P. albipes* and *P. cora* in New Zealand, were rather unusual. They made their nests almost exclusively with *Deleatidium* mayflies, female wasps wait on rocks in the stream until they encounter a *Deleatidium* nymph in the process of molting to the subimago (Harries, 1990). A muscid fly *Spilogona* sp. completed with the wasps for subimagos emergence on the rocks in the stream. Similarly, the larvae of therevid fly *Anabarynychus* sp. burrows through the soil

along the edge of the stream. When this larva encountered a provisioned nest *Podagritis*, it consumed the mayfly *Deleatidium* (Harries, 1990).

According to Dudgeon (1989) the nymph of damselfly, *Euphaea decorata* has showed preference toward the nymphs of mayflies by increase of age. The number of genera for predation was twelve and high percent of the prey consumed by small nymphs was for two genera of mayflies, while a large nymphs fed on five genera which represented 36% of the prey consumed. According to Rabeni (1992) mayflies were the larger portion of the food for young smallmouth bass, *Micropterus dotomieu* and rock bass, *Ambloplites rupestris* with still larger portion of the food for older white suckers, *Catostomus commersoni* (Chen and Harvey, 1995) and common river galaxias, *Galaxias vulgaris* (Cadwallader 1975). The change in the food preference of galaxias occurred when it moved from quiet water to riffles have a greater diversity of food items (Cadwallader 1975).

According to Caira and Orringer (1995), fishes feed predominantly on mayflies. Young freshwater stingrays, *Potamotrygon magdalenae*, feed exclusively on



mayflies. The relative frequency of mayflies in the gut of some fishes was very high; some species exhibited a wide range of preference to young small mouth bass, *Micropterus dolomieu* (Easton *et al.*, 1996), Bluegilled bully, *Gobiomorphe hubbasi*, and torrentfish, *Cheimarrichthus fosteri*, green sandpipers, *Tringa ochropus*, green bee-eater, *Merops orientalis*, *Chiroptera*, *Pterops giganteus* (Scrimgeour and Winterbourn, 1987). Salamander (Gulf coast waterdog) *Necturus beyeri*, preferred *Leptophelebia* mayflies to younger and *Stenonema* mayflies to older forms (Bart and Holzenthal, 1985). The birds, *Eurasian dipper*, *Cinclus cinclus* consumed more mayflies during the breeding season (Cormerod and Tyler, 1991) as like the black duck, *Anus rubripes* (Reinecke and Owen, 1980). Similarly, the salamander, *Leurognathus marmorata* found feeding on 10 species of Mayflies (Mart-of and Scott 1957).

The Pyrenean desman *Galemys pyrenaicus* (Mammalia) found feeding commonly on mayflies, about 96% of all the guts examined contained mayflies body parts (Castien and Gosalbez 1995). However, these mayflies only represented 16% of the food ingested by volume. According to Castien and Gosalbez (1995), mayflies do not account for a large volume of the food consumed. However, mayflies by small predators abundant in the stream easily captured and consumed. According to Reinecke and Owen (1980), mayflies represented 29% of the grass energy ingested in black ducks. Similarly, the green sandpiper, *Tringa ochropus* consumed 9,500 to 11,000 *Baetis* nymph per day (Ormerod and Tyler, 1988). Mayflies are also recognized as human (*Homo sapiens*) diet (Bodenheimer, 1951) described. In North Vietnam, China and Japan mayflies largely consumed by humans. The people of Malawi make a paste out of Mayflies (*Caenis kungu*) and mosquito called kungu (Fladung, 1924).

Gut content analysis of nymphs by Nost (1985) showed to be herbivore- detritovore and that there was a seasonal shift in the food items found in the gut from green algae during December to April to detritus and to some extent, flagellates, during June to July. From June till around October, dense algal mats were observed to cover the *Hydrilla* strands, a microscopic examination of which revealed the predominance of *Spirogyra*, *Oedogonium*, *Closterium*, *Cosmarium*, *Staurostrum*, *Oscillatoria* and *Mougeotia* (Gupta *et al.*, 1993) and these were forms which constituted a major portion of the food of nymphs during this period. Several studies reveal that algae were the food of superior quality than detritus (Cummins & Klug, 1979, Bird and Kaushik, 1984). The observations of on growth of nymphs fed algae/detritus also showed that mayflies grow and reach maturity faster on an algal diet (Gupta *et al.*, 1993). Thus, it seems probable that several factors such as a higher temperature regime, abundance of energy rich algal food and probably reduced predation pressure, were responsible for the population buildup of nymphs during March to June, although studies that are more

specific would have to be conducted to understand further the role of these factors in predation.

Okedi (1992) noted that the traditional protein for some domestic African animals including fish, have declined over the years. He considered using the lakeflies as a source of protein. His analysis along with that of Bergeon *et al.*, (1988) show that these insects are high in protein, minerals, B vitamins and essential amino acids. They were also low in fat and moisture and thus contributed for long life. The insect's cakes have a high digestibility. *Povilla* occurred in huge swarms made them relatively easy to capture and to use as food. However, suitable commercial harvesting technique needs to be designed. Very recently, Sathe (2015) exposed some insects as human diet including their preparation and values.

## Acknowledgment

Authors are thankful to Shivaji University Kolhapur and BARTI, Pune for Providing facilities and financial assistance to Smt. Rohini Kamble.

## Conflict of Interests

Authors declare that there is no conflict of interests regarding the publication of this paper.

## References

- [1]. Anderson, N.H., and K.W. Cummins. Influences of diet on the life histories of aquatic insects *J. Fish. Res. Board Can.* 36: 335-342 (1979).
- [2]. Bart, H.L., and R. W. Holzenthal. Feeding ecology *Necturus beyeri* in Louisiana. *J. Herpetol.* 19: 402-410 (1985).
- [3]. Bergeron, D., R.J. Bushway, F.L. Roberts, I Kornfield, J. Okedi and A.A. Bushway. The nutrient composition of an insect's flour sample from Lake Victoria, Uganda. *J. Food Comp. Anal.* 1: 371-377 (1988).
- [4]. Bodenheimer, F. S. Insects as human food. A chapter of the ecology of *Man. W. Junk, Publisher, The Hague* (1951).
- [5]. Brittain, J.E. Semivoltinism in mountain populations of *Nemurella pictetii* (Pecoptera). *Oikos* 29:302-305 (1978).
- [6]. Caira, J. N and D.J. Orringer. Additional information on the morphology of *Potamotrygonocetus magdalensis* (Tetraphylidae : Onchobothriidae) from the freshwater stingery *Potamotrygon Magdalena's* in Colombia. *J. Helminthol. Soc.* 62: 22.26 (1995).
- [7]. Castien, E. and J. Gosalbez. Diet of *Galemys pyrenaicus* (Geoffroy, 1811) in the north of the Iberian Peninsula. Netherlands. *J. Zool.* 45: 422-430 (1995).

- [8]. Cummins, K.W., R. C. Petersen, F.O.Howard, J.c.Wuychck, and V.I.Holt. The utilization of leaf litter by stream detritivores. *Ecology* 54: 336-345 (1973).
- [9]. Downes, J. A. Feeding and mating in the insectivorous ceratopogoninae (Diptera). *Mam. Ent Soc. Can.* 104: 1-62 (1978).
- [10]. Dudgeon. D. Life cycle, Production, microdistribution and diet of the damselfly *Euphaea decorata* (Odonta: Euphaeide) in a Hong Kong Forest stream. *J. Zool.* 217: 57-72 (1989).
- [11]. Fladung, E. B. Insect as food. *Maryland Acad. Sci. Bull.* 5-8 (1924).
- [12]. Gibbs, K.E. and T. M. Mingo. The life history, nymphal growth rates, and feeding habits of *Siphonisca aerodromia* Needham (Ephemeroptera: Siphonuridae) in Marine. *Can. J. Zool.* 64: 427-430 (1986).
- [13]. Giller, P.S. The natural diet of the Notonectidae: Field trials using electrophoresis. *Ecol. Ent.* 11: 163-172 (1986).
- [14]. Gupta, S.J., Michael, R.G and Gupta, A. Influence of diet on growth, food retention time, and gill ventilation rate of nymphs of *cloeon* sp. (Ephemeroptera : Baetidae). *Hydrobiologia* (1993).
- [15]. Harries, A.C. *Podagritys cora* (Cameron) and *P. albines* (F. Smith) (Hymenoptera: Specidae: Cabroninae) preying on Ephemeroptera and Trichoptera, *Pan-Pac. Ent.* 66: 55-61 (1990).
- [16]. Kamble Rohini and Sathe T.V. Diversity and Economic importance. *Indian Jornal of Applied Reseaech* 5: 7, 2249-555 (2015).
- [17]. Kamble Rohini and Sathe T.V. Biology and Morphology of Nymph of *Indialis badia* (Ephemeroptera: Leptophelebiidae). *Indian Jornal of Applied Reseaech* 6: 5, 74.50 (2016).
- [18]. Kjellberg, G. Autekologiska studies over *Leptophelebia vespertina* (Ephemeroptera) i en mindre skogstiarn (1966-1968). *Ent. Tidskar*, 93:1-29 (1992).
- [19]. Lau DCP, Leung KMY, Dudgeon D. Experimental dietary manipulations for determining the relative importance of allochthonous and autochthonous
- [20]. food resources in tropical streams. *Freshwater Biol*, 53:139–147 (2008).
- [21]. Lau DCP, Leung KMY, Dudgeon D. Are autochthonous foods more
- [22]. Important than allochthonous resources to benthic consumers in tropical
- [23]. headwater streams? *J N Am Benthol Soc*, 28:426–439 (2009).
- [24]. Li AOY, Dudgeon D. Food resources of shredders and other benthic
- [25]. Macroinvertebrates across a range of shading conditions in tropical Hong
- [26]. Kong streams. *Freshwater Biol*, 53:2011–2025 (2008).
- [27]. Mantel SK, Salas M, Dudgeon D. Food web structure in a tropical Asian
- [28]. forest stream. *J N Am Benthol Soc*, 23:728–755 (2004).
- [29]. March JG, Pringle CM. Food web structure and basal resource utilization
- [30]. along a tropical island stream continuum, Puerto Rico. *Biotropica*, 35:84–93 (2003).
- [31]. Martof, B.s and D.C.Scott. The food of the salamander *Leurognathus*. *Ecology*, 38: 495-501 (1957).
- [32]. Nost, T. Distribution and food habits of Mayflies (Ephemeroptera) in streams in the Dovrefjell Mountains. Central Norway. *Fauna Norv. Ser. B.* 32:100-105 (1985).
- [33]. Ormerod, S.J and tyler. Exploitation of prey by a river bird, the dipper *Cinclus cinclus* (L.), along acidic and circumneutral streams in upland Wales. *Freshwater Biol*, 25: 105-116 (1991).
- [34]. Rabeni, C.F. Trophic linkage between stream centrachids and their crayfish prey. *Cn J.Fish. Aquat. Sci*, 49: 1714-1721 (1992).
- [35]. Reinecke, K.J. and R.B.Owen, Jr. Food use and nutrition of black ducks nesting in Marine. *J. Wildl. Manage.* 44: 549-558 (1980).
- [36]. Sathe T.V. Recents trends in biological pest control. Astral Int. Nat. Pvt. Ltd, New Delhi. pp 1-204 (2014).
- [37]. Sathe T.V. Insect for human diet from Kolhapur region, India. *Int. J. Pharm. Bio. Sci.* 6(1): (B) 519-527 (2015).
- [38]. Sateesh Pujari, & Estari Mamidala. (2015). Anti-diabetic activity of Physagulin-F isolated from *Physalis angulata* fruits. *The American Journal of Science and Medical Research*, 1(2), 53–60. <https://doi.org/10.5281/zenodo.7352308>
- [39]. Sathe T. V. & R.N. Patil. Conservation protection and utilization strategies for Amphibians from Satara district Maharashtra including Ghats. *Biolife*, 4(1): 189-194 (2016).
- [40]. Sweeney. B.W. factors influencing life history pattern of aquatic insects. In *Ecology of aquatic insect* (Eq.V.H.Resh and D Rosenberg), pp 56-100. Preager scientific publishers, New York (1984).
- [41]. Scrimgeour, G.j and M.J.Winterbourn. Diet, food resource partitioning and feeding periodicity of two riffle-dwelling fish species in a New Zealand river. *J.Fish.Biol.* 31: 309-324 (1987).
- [42]. Soluk, D.A. and H.F. Clifford. Life history and abundane of the predaceous psammophilous mayfly *Pseudiron centrali* McDunnough (Ephemeroptera: Heptageniidae). *Can. J.Zool.* 62: 1534-1539 (1984).
- [43]. Wallance, J. B., W.R. Woodali., and F.F.Sherberger. Breckdown of leaves by feeding of *Peltoperla maria* Nymph (Plecoptera : Peltoperlidae). *Ann. ent. Soc. Am.* 63: 562-567 (1970).